Table 4

Cancer Incidence in Wayland, MA 1987-1994

1987-1994	Total					Male					Female							
	Obs	Exp	SIR	95	%	CI	Obs	Exp	SIR	95	%	CI	Obs	Exp	SIR	95	%	CI
Bladder	14	17.1	82	45		137	10	12.8	78	38		144	4	4.3	NC		NC	
Breast	82	75.2	109	87		135	1	0.6	NC		NC		81	74.7	108	86		135
Hodgkin's Disease	4	3.5	NC		NC		1	1.9	NC		NC		3	1.6	NC		NC	
Kidney	11	11.6	95	47		170	8	7.4	109	47		214	3	4.2	NC		NC	
Leukemia	6	8.1	74	27		161	3	4.8	NC		NC		3	3.3	NC		NC	
Liver	3	2.3	NC		NC		2	1.8	NC		NC		1	0.6	NC		NC	
Lung #	63	70.1	90	69		115	32	43.2	74	51		104	31	26.9	115	78		164
NHL	18	16.1	112	66		177	7	8.8	80	32		164	11	7.3	137	75		270
Pancreas	6	9.0	67	24		145	5	4.7	107	35		251	1	4.3	NC		NC	
Stomach	5	10.0	50	16		116	4	6.9	NC		NC		1	3.1	NC		NC	

Notes:

Expected number of cases presented are rounded to the nearest tenth.

SIRs are calculated based on the exact number of expected cases.

SIRs and 95% Confidence Interval are not calculated when fewer than 5 cases are observed.

Obs = Observed number of cases

Exp = Expected number of cases

SIR = Standardized Incidence Ratio

NC = Not Calculated

95% CI = 95% Confidence Interval

* Indicates statistical significance (P<.05)

One case of lung cancer could not be mapped due to insufficient address information and was not assigned to a census tract. This case was included in the town totals.

Data source: Massachusetts Cancer Registry, Bureau of Health Statistics, Research and Evaluation, Massachusetts Department of Public Health Data analysis: Community Assessment Unit, Bureau of Environmental Health Assessment, Massachusetts Department of Public Health